

# Atlas of Genetics and Cytogenetics in Oncology and Haematology

OPEN ACCESS JOURNAL



INIST-CNRS

## Gene Section Review

### SLC6A4 (solute carrier family 6 member 4)

Rafiq Gurbanov, Berkay Kalkanci

Department of Molecular Biology and Genetics (RG); Bilecik Seyh Edebali University, Department of Biotechnology (BK), Bilecik Seyh Edebali University, Biotechnology Application and Research Center, Bilecik, Turkey; rafiq.gurbanov@bilecik.edu.tr; berkaykalkancii@gmail.com

Published in Atlas Database: March 2019

Online updated version : <http://AtlasGeneticsOncology.org/Genes/SLC6A4ID49951ch17q11.html>

Printable original version : <http://documents.irevues.inist.fr/bitstream/handle/2042/70648/03-2019-SLC6A4ID49951ch17q11.pdf>

DOI: 10.4267/2042/70648

This work is licensed under a Creative Commons Attribution-Noncommercial-No Derivative Works 2.0 France Licence.

© 2020 Atlas of Genetics and Cytogenetics in Oncology and Haematology

### Abstract

The SLC6A4 gene encodes a sodium-dependent serotonin reuptake protein delivering the neurotransmitter serotonin from the synaptic cleft back to the presynaptic end. Its main function is to abort the activity of serotonin and forward it to neurotransmitter pool for recycling. The psychomotor stimulant drugs mainly amphetamines and cocaine act on this transmembrane protein which is a member of the sodium: neurotransmitter symporter family. SLC6A4 gene polymorphisms affect the rate of serotonin reuptake and play an important role in pathogenesis of various illnesses like Sudden infant death syndrome, aggressive

behaviour in Alzheimer patients, Seasonal affective disorder, Major depressive Disorder and Obsessive-compulsive disorder.

### Keywords


Solute Carrier Family 6 Member 4 (SLC6A4), Serotonin (5-HT), Serotonin transporter protein (5-HTT), Anxiety, Alcoholism, Hypertension, Mental Disorders.

### Identity


**Other names:** 5-HTT, 5-HTTLPR, 5HTT, HTT, OCD1, SERT, SERT1, hSERT

**HGNC (Hugo):** SLC6A4


**Location:** 17q11.2



**Figure 1.** Genomic location of SLC6A4 (Chromosome 17 - NC\_000017.11 Reference GRCh38.p12 Primary Assembly)



**Figure 2. Chromosomal Location of SLC6A4.** Cytogetic Location: 17q11.2, which is the long (q) arm of chromosome 17 at position 11.2 (U.S. National Library of Medicine)



**Figure 3.** Numbers and illustrative sizes of exons of human SLC6A4. (<https://www.atlasgeneticsoncology.org>)

## DNA/RNA

The SLC6A4 gene is 41,684 bp long (according to UCSC, GRCh38/hg38), located on the plus strand and spans 15 exons (NCBI Homo sapiens Annotation Release 109).

### Transcription

The gene has 5 transcripts (Table 1)

Name	Transcript ID	bp	Protein (aa)	Biotype
SLC6A4-201	ENST00000261707.7	6604	630 aa	Protein coding
SLC6A4-202	ENST00000394821.2	2160	618 aa	Protein coding
SLC6A4-203	ENST00000401766.6	6543	630 aa	Protein coding
SLC6A4-204	ENST00000578609.1	566	No protein	Retained intron
SLC6A4-205	ENST00000579221.5	1069	72 aa	Nonsense mediated decay


Table 1. Transcripts of human SLC6A4 gene (Ensemble, GRCh38.p12).

## Protein

SLC6A4 gene encodes a serotonin transporter protein (5-HTT) of 70,320 Dalton and composed of 630 amino acids (Figure 4) (Ramamoorthy et al., 1993). The protein belongs to family of neurotransmitter/sodium transporter (NSS). NSS family also includes dopamine, glycine and  $\gamma$ -aminobutyric acid (GABA) transporters (Chen et al., 2004). The members of this family have 12 transmembrane domains and intracellular N and C terminal regions (Yamashita et al., 2005). The large extracellular structure (EL) between TM3 and TM4 is modified by N-linked glycosylation in all eukaryotic NSS proteins and the number of N-linked glycosylation sites can vary from carrier to carrier. The consensus sequence for N-linked glycosylation

is N-X-S/T (the amide nitrogen glycan binding region in the asparagine side chain and X is any amino acid except proline) (Mitra et al., 2006).

The EL2 of the 5-HTT protein has two glycosylation sites carrying glycan (Tate and Blakely, 1994). Mutations in the glycosylation sites of 5-HTT and other NSS proteins generally cause problems in the transport of neurotransmitter proteins such as serotonin at the cell surface (Tate and Blakely, 1994; Olivares et al., 1995; Melikian et al., 1996; Li et al., 2004).



**Figure 4.** Structure of human Solute carrier family 6 member 4. X-ray structure of the ts3 human serotonin transporter complexed with paroxetine (selective serotonin re-uptake inhibitor) at the central site (PDB ID: 5L6X) (<https://www.rcsb.org>).

### Expression

SLC6A4 gene is most commonly expressed in gastrointestinal tract, female tissues, and lung. It is less expressed in muscle, skin, endocrine, male, and female tissues (<http://www.proteinatlas.org>).


### Localisation

SLC6A4 is found in various cellular compartments such as cytosol, endosome, plasma membrane, and integral component of plasma membrane, integral component of postsynaptic membrane, integral component of presynaptic membrane, endomembrane system, neuron projection, and serotonergic synapse (Müller et al., 2006; Brenner et

al., 2007; Ahmed et al., 2008; Ahmed et al., 2009; Gaudet et al., 2011).

### Function

5-HTT protein activity relies on the concentrations of intracellular potassium and extracellular sodium and chloride ions. It also depends on the membrane potential generated by sodium-potassium adenosine triphosphatase for the activity of the 5-HTT protein. The 5-HTT protein binds itself to sodium, serotonin and chloride ions, respectively.



**Figure 5.** Expression profile of SLC6A4 in different tissues/organs in human. Data were taken from The Human Protein Atlas (<http://www.proteinatlas.org>) in December, 2018.

Thus, the membrane potential mediates the release of sodium and chloride molecules pre-bound to the 5-HTT protein and the 5-HTT protein passes into the cell. The 5-HTT protein releases serotonin in the cell and binds a potassium ion to itself. 5-HTT is activated by potassium ion and may be out of the cell. Serotonin (5-hydroxytryptamine; 5-HT) is an important neurotransmitter substance in the central and peripheral nervous system. After release of 5-HT into brain synapses, the  $\text{Na}^+$  and  $\text{Cl}^-$  ions-dependent high-affinity serotonin transporter SLC6A4 (also called 5-hydroxytryptamine transporter, 5-HTT, SERT), which is localized in presynaptic neuronal membranes, effectively clears 5-HT from the synaptic space. Thus, the synaptic activity of 5-HT is terminated by 5-HTT and reintroduced into the neurotransmitter pool for re-use. In this way, 5-HTT has an important role in the retrieval of serotonin and

the execution of serotonergic function (Gelernter J. Et al., 1998, Catalano M., 1999).

The 5-HTT activity is rapidly regulated by a number of G-protein-linked receptors and protein kinase-associated pathways including protein kinase C (PKC), PRKG1 (protein kinase G, PKG), and p38 mitogen-activated protein kinase (MAPK). The PKC-dependent phosphorylation and down-regulation of the 5-HTT protein is sensitive to extracellular 5-HT and plays a regulatory role in the transport of 5-HT (Ramamoorthy et al., 1998; Ramamoorthy et al., 1999; Zhu et al., 2004; Samuvel et al., 2005; Prasad et al., 2005).

### Mutations


Most frequent mutations are located in the 5-HTTLPR of the promoter region of SLC6A4 gene. These mutations lead to the clinical picture of Autism (rs6365, rs28914832, rs140700) (Sutcliffe et al., 2005; Landaas et al., 2010; Adamsen et al., 2011), Increased rigid-compulsive behaviour in autism (rs28914833, rs28914834) (Sutcliffe et al., 2005; Rao et al., 2017), Obsessive-compulsive disorder (rs25532) (Wendland et al., 2007), Major depressive disorder (rs6354) (Rao et al., 2017), Panic disorder (rs3813034) (Gyawali et al., 2010), Unipolar disorder (Ogilvie et al., 1996), and Pulmonary arterial hypertension (Eddahibi et al., 2003) (Table 2).

### Implicated in

The serotonin carrier protein encoded by SLC6A4 gene is the target of serotonin selective reuptake inhibitors, an important class of antidepressant drugs (Ramos et al., 2006). Three alleles of 17-bp VNTR (variable number tandem repeat) were detected in the intron 2 region of the gene, between 9 (Stin2.9), 10 (Stin2.10), and 12 (Stin2.12) copies. Presence of Stin2.9 allele in humans has been reported to increase the risk of Major depressive disorder (MDD) (Ogilvie et al., 1996).

The 5-HTTLPR promoter sequence of the gene comprises two variant repeat length polymorphisms, known as the 16-element long (L) and 44-bp short (S) variant with 14 repetitive elements (Esterling et al., 1998; Sen et al., 2004). The L/L genotype from the 5-HTTLPR promoter variants causes more 5-HTT protein expression than the L/S or S/S variants (Canli and Lesch, 2007).

Analysis of lymphoblastoid cell lines with different genotypes revealed that the basal activity of the L variant of the SLC6A4 gene promoter was two times greater than that of the S variant (Lesch et al., 1996).



**Figure 6.** Serotonergic Synapse Pathway Map. Once released from presynaptic axonal terminals, 5-HT binds to receptors, which have been divided into 7 subfamilies on the basis of conserved structures and signalling mechanisms. These families include the ionotropic 5-HT3 receptors and G-protein-coupled 5-HT receptors, the 5-HT1 (Gi /Go -coupled), 5-HT2 (Gq-coupled), 5-HT4/6/7 (Gs-coupled) and 5-HT5 receptors. Presynaptically localized 5-HT1B receptors are thought to be the autoreceptors that suppress excess 5-HT release. 5-HT's actions are terminated by transporter- mediated reuptake into neurons, leading to catabolism by monoamine oxidase. Data were taken from KEGG: Kyoto Encyclopedia of Genes and Genomes (<https://www.genome.jp/kegg/>) in February, 2019.

Pairwise Alignment Scores

Gene		Identity (%)	
Species	Symbol	Protein	DNA
<b>H.sapiens</b>		<b>SLC6A4</b>	
vs. P.troglodytes	SLC6A4	99.4	99.3
vs. M.mulatta	SLC6A4	98.6	97.5
vs. C.lupus	SLC6A4	93.5	90.3
vs. B.taurus	SLC6A4	93.0	89.7
vs. M.musculus	Slc6a4	92.5	87.8
vs. R.norvegicus	Slc6a4	91.7	88.1
vs. G.gallus	SLC6A4	81.9	76.9
vs. X.tropicalis	LOC100493853	76.8	71.1
vs. D.rerio	slc6a4a	70.8	68.3

**Figure 7.** Pairwise alignment of SLC6A4 gene protein sequences (in distance from human) (HomoloGene, NCBI).

Missense/Nonsense Mutations					
#	Location	Mutation	Protein	Reported Phenotype	Reference
1	Exon 2	c.10A>G	p.T4A	Enhanced 5-HT transport activity	Parasad et al., 2005
2	Exon2	c.167G>C	p.G56A	Autism, association with	Sutcliffe et al., 2005 Camarena et al., 2012
3	chr17:30218213	c.603G>C	p.K201N	Increased transporter activity	Rasmussen et al., 2009
4	Exon 4	c.643G>A	p.E215K	MAPK nonresponsiveness	Parasad et al., 2005
5	Exon 6	c.878C>T	p.S293F	Enhanced 5-HT transport activity	Parasad et al., 2005
6	Exon 7	c.1016C>T	p.P339L	Reduced uptake activity	Parasad et al., 2005
7	Exon 8	c.1084C>A	p.L362M	Enhanced 5-HT transport activity	Parasad et al., 2005
8	Exon 9	c.1273A>C	p.I425L	Autism, association with	Sutcliffe et al., 2005
9	Exon 9	c.1273A>G	p.I425V	Obsessive-compulsive disorder, susceptibility, association	Ozaki et al., 2003 Moya et al., 2013
10	Exon 10	c.1393T>C	p.F465L	Increased rigid-compulsive behavior in autism, association with	Sutcliffe et al., 2005
11	Exon 12	c.1648C>G	p.L550V	Increased rigid-compulsive behavior in autism, association with	Sutcliffe et al., 2005 Rao et al., 2017
12	Exon13	c.1815A>C	p.K605N	MAPK nonresponsiveness	Parasad et al., 2005 Rao et al., 2017
13	Exon 14	c.1861C>T	p.P621S	MAPK nonresponsiveness	Parasad et al., 2005
Splicing Mutation					
14	chr17:30216371	c.838-155G>A	NA	Autism, association with	Sutcliffe et al., 2005 Landaas et al., 2010
Regulatory Mutation					
15	chr17:30237328	c.-1936G>A	NA	Obsessive-compulsive disorder, association with	Hu et al., 2006 Perroud et al., 2010 Hung et al., 2011 Moya et al., 2013 Stacey et al., 2013
16	chr17:30237152	c.-1760T>C	NA	Obsessive-compulsive disorder, association with	Wendland et al., 2008 Fuxman Bass et al., 2015
17	chr17:30222880	c.-185A>C	NA	Major depressive disorder, association with	Rao et al., 2017
18	chr17:30197993	c.463T>G	NA	Increased expression	Vallender et al., 2008

					Seneviratne et al., 2009
19	chr17:30197786	c.670T>G	Na	Panic disorder, association with	Gyawali et al., 2010 Aoki et al., 2010 Hartley et al., 2012
<b>Deletion</b>					
20	Promoter	c.1212-1255 del TGCAGCC	NA	Anxiety related traits, association with	Helis et al., 1996 Marziniak et al., 2005 Borroni et al., 2006 Albani et al., 2009
	<b>Location</b>	<b>Repetition Sequence</b>	<b>Number Of Repetitions</b>	<b>Reported Disease/Phenotype</b>	<b>Reference</b>
21	Intron 2	(GGCTGYGACCYRGRRTG) <sub>n</sub>	10-12	Unipolar disorder, association with	Ogilvie et al., 1996
22	Intron 2	(GGCTGYGACCYRGRRTG) <sub>n</sub>	12	Pulmonary arterial hypertension, association with	Allen et al., 2008: Cao et al., 2009

Table 2. Solute Carrier Family 6 Member 4 (SLC6A4) related mutations.

In a later study, the expression of the native SLC6A4 gene in cultured lymphoblast cell lines from subjects with different SLC6A4 promoter genotypes was examined (Lesch et al., 1996). The mRNA concentration of the SLC6A4 gene in L/L cells was found to be 1.4 to 1.7-fold higher than the L/S and S/S cells (Lesch et al., 1996). Bradley et al. (2005) directly measured serotonin transporter mRNA levels and also identified 4 loci containing the serotonin transporter gene from 85 independent lymphoblast lines. They found strong impact of 5-HTTLPR on the mRNA expression (Bradley et al., 2005).

The Gly56Ala substitution in the exon 2 of the SLC6A4 gene has been reported to be associated with autism and exhibit structurally high SERT activity (Sutcliffe et al., 2005). I425V substitution in the exon 9 of the SLC6A4 gene has been reported to be associated with obsessive-compulsive disorder (OCD) (Ozaki et al., 2003; Kilic et al., 2003; Zhang et al., 2007). In addition, A-1438G and T102C polymorphisms were reported to be associated with OCD (Taylor, 2013; Taylor, 2016). Rao et al. (2017) reported that I550V (exon 12) and K605N (exon 13) substitutions of the SLC6A4 gene are associated with major depression disorder (MDD) and SA (non-fatal suicidal behavior) in addition to autism and OCD in 36 Chinese patients.

#### Animal Experiments

When 5-HTT was temporarily inhibited by fluoxetine (selective serotonin re-uptake inhibitor) in

the early developmental period of mice, it was observed that adult mice exhibited abnormal emotional behaviors. It has been reported that serotonin plays a critical role in the maturation of brain systems that regulate emotional function, and that the low expression level of the SLC6A4 gene may be related to the development of psychiatric disorders in adults (Ansorge et al., 2004). A study of transgenic mice overexpressing the SLC6A4 gene and wild-type mice showed that right ventricular pressure was 3-fold higher in transgenic mice (Maclean et al., 2004). Page et al. (2009) reported macrocephaly in Pten +/- mice. Female Pten +/- mice had socialization disorder, whereas male Pten +/- mice had no socialization disorder. This phenotype was exacerbated in mice with Pten and SLC6A4 double haploinsufficiency. As a result of these findings PTEN and SLC6A4 genes have been reported to be associated with autism spectrum disorder (ASD).

#### Breast cancer

High expression is found in breast cancer, but the gene product is not prognostic according to The Human Protein Atlas.

A fusion gene PIP4K2B /SLC6A4 was found in breast cancer (Yoshida et al., 2013).



### **Obsessive-Compulsive Disorder (OCD)**

Hu et al. (2006) found that the gain-of-function homozygous L(A)L(A) genotype was 2-fold in patients with OCD compared to healthy individuals. In a replication study in 175 trios consisting of probands with OCD and their parents, the L(A) allele was 2-fold overtransmitted to the patients with OCD. The HTTLPR L(A)L(A) genotype exerted 1.8-fold effect on risk of OCD, thus establishing the role of the HTT gene in OCD. In another study, the frequency of Stin2.12 allele in Asian patients with anxiety disorder (including OCD) was reported to be significantly higher compared to the healthy individuals (Ohara et al., 1998). There is also a possible association between OCD and Stin2.12 allele in Spanish Caucasian population (Baca-Garcia et al., 2007; Saiz et al., 2008).

### **Anxiety-Related Personality Traits**

The homozygous or heterozygous form of the S-variant of the 5-HTTLPR polymorphism in the SLC6A4 gene has been reported to be associated with lower expression and openness, and higher neuroticism (Lesch et al., 1996). Individuals with 1-2 copies of the S-variant of the 5-HTTLPR polymorphism, which is implicated in reduced 5-HTT expression and function and increased fear and anxiety-related behaviours, demonstrated higher amygdala activity in response to fearful stimuli compared with individuals homozygous for the L-variant (Hariri et al., 2002). The altered function of the serotonin neurotransmission system causes aggressive behaviour in Alzheimer's patients (AD) (Brown et al., 1982; Palmer et al., 1988). In a study conducted on 137 AD patients, the aggressive behaviours (58 patients) were associated with L/L genotype (Sukonick et al., 2001).

### **Major Depressive Disorder (MDD)**

People with L/S or S/S allele in stressful life conditions have been reported to have more depression and suicidal tendency than those with L/L allele (Caspi et al., 2003). They also reported that the individual's response to environmental insults was determined by environmental-gene interaction (Caspi et al., 2003). In a study conducted on Pomerania population, the relationship of S-variant with environmental interaction and depression was determined. The results of the study supported previous studies on the gene-environment interaction of the S allele. It has also been revealed to cause high mental vulnerability to social stress and chronic diseases (Grabe et al., 2005). Homozygous or heterozygous genotypes of the S-variant were associated with stress-related depression and anxiety disorders in elite athletes (Petito et al., 2016). The 5-HTTLPR polymorphisms of the SLC6A4 gene have been reported to be associated with prenatal and

postnatal depression (Sanjuan et al., 2008; Oberlander et al., 2013). Interestingly, the genetic and epigenetic variations have been reported in the SLC6A4 gene of children exposed to prenatal depression (Devlin et al., 2010; Oppenheimer et al., 2013; Wankerl et al., 2014; Babineau et al., 2014; Green et al., 2017). 5-HTTLPR polymorphisms and DNA methylations of SLC6A4 gene have been reported to be associated with depression (Devlin et al., 2010; Sugawara et al., 2013).

### **Seasonal Affective Disorder (SAD)**

Willeit et al. (2003) genotyped 284 subjects (138 SAD patients and 146 healthy individuals) to examine the relationship between L and S-variants in the 5-HTTLPR polymorphism of SLC6A4 gene with SAD. The distribution of genotype and S allele frequency was found similar between patients and healthy subjects, while they were correlated with the subtypes of DSM-IV depression. L allele was correlated with melancholic depression whereas S allele was related to atypical depression. It was concluded that 5-HTTLPR polymorphism affects the phenotypic disease expression but it is not the cause of disease.

The effects of light therapy on serotonin transporter binding (5-HTT BP<sub>ND</sub>), which is biomarker of 5-HTT levels, in the anterior cingulate and prefrontal cortices (ACC and PFC) of healthy individuals during the fall and winter was studied. In winter, light therapy significantly decreased 5-HTT BP<sub>ND</sub> by 12% in the ACC with respect to placebo, whereas in the fall, no significant change in 5-HTT BP<sub>ND</sub> was measured. In this context, it has been reported that 5-HTT BP<sub>ND</sub> can be used as a biomarker for the assessment of the modification effects of light therapy (Harrison et al., 2015). In a study on 20 SAD patients and 20 healthy participants the impact of seasonal (winter and summer) variations on 5-HTT activity was investigated by analysing brain 5-HTT BP<sub>ND</sub> levels. The study reported a significant increase in 5-HTT BP<sub>ND</sub> in different brain regions (including ACC and PFC) especially in severe SAD during winter. The 5-HTT BP<sub>ND</sub> as biomarker in the diagnosis of SAD is important as it can be applied for the development of prevention strategies against disease progress (Tyrer et al., 2016).

### **Alcoholism**

A meta-analysis study (data from 3,489 alcoholics and 2,325 controls) was conducted on the association of 5-HTTLPR polymorphisms of SLC6A4 with alcohol dependence. It has been reported that alcohol dependence complicated by a comorbid psychiatric condition or a more severe form of alcoholism is associated with the S-variant (Feinn et al., 2005). In a study conducted on university students, the S-variant was reported to be associated with high alcohol consumption (Herman et al., 2003). High

alcohol consumption in men with homozygous or heterozygote genotypes of the S-variant has been reported, whereas it was seen only in the heterozygous women (Munaf et al., 2005). In a study conducted on 273 (78.5% male) alcoholic individuals of the Caucasus and Hispanic origin, it was reported that G allele carriers for the rs1042173 SNP in the 3'UTR region of the SLC6A4 gene had lower alcohol consumption than T-allele homozygotes (Seneviratne et al., 2009). G allele transfection to HeLa cells resulted in higher mRNA and protein expression compared to T allele transfected cells (Seneviratne et al., 2009).

### **Sudden Infant Death Syndrome (SIDS)**

L/L genotype and excess of L-variant have been reported to be associated with SIDS (Narita et al., 2001; Weese-Mayer et al., 2003). In addition, the Stin2.12 allele has been associated with SIDS in African-Americans and Japanese, whereas it is not associated with SIDS in Caucasian-Americans (Narita et al., 2001; Weese-Mayer et al., 2003).

### **Panic Disorder (PD)**

The rs3813034 variant in the 3'UTR region of the SLC6A4 gene has been reported to be related to PD (Gyawali et al., 2010). Decreased SLC6A4 gene expression due to the gene polymorphisms in midbrain, hypothalamus and temporal lobe has been associated with PD (Maron et al., 2006). Strug et al. (2010) reported that rs140701 variant of the SLC6A4 gene may be associated with PD.

### **Bipolar Affective Disorder (BPAD) or Manic-Depressive Illness (MDI)**

The disorder of the neurotransmitter system, including serotonin and monoamines, has been reported to be associated with bipolar disorder (Scott et al., 1979; Kapur and Remington, 1992). The patients with BPAD without panic disorder exhibited statistically higher frequencies of the Catechol O-Methyltransferase (COMT) Met158 and the S-variant 5-HTTLPR genotypes with respect to healthy individuals (Rotondo et al., 2002). When the relationship between L and S-variant polymorphisms of SLC6A4 gene and BPAD and unipolar depression were investigated by meta-analysis study, it was revealed that the S-variant could be associated with BPAD but not with unipolar depression. The L-variant was not implicated in BPAD and/or unipolar depression (Lasky-Su et al., 2005). However other studies reported no significant relationship between SLC6A4 5-HTTLPR and VNTR polymorphisms and BPAD (Mendlewicz et al., 2004; Cho et al., 2005). The association between epigenetic variations of SLC6A4 gene and BPAD were studied on 20 bipolar monozygotic twins with respect to 20 healthy subjects. The promoter-wide

DNA methylation analysis of lymphoblastic cell lines (LCLs) revealed DNA hypermethylation in SLC6A4 gene that can be an epigenetic mark resulting from a G×E interaction leading to the development of BPED (Sugawara et al., 2011). In a study on the association of SLC6A4 and DRD2 (dopamine D2 receptor) genes with BPED, the gender specific differences in SLC6A4 gene polymorphisms were reported. The results revealed gender-specific association of the DRD2 A1/A1 and the 5-HTTLPR S/S, S/LG, and LG/LG (S+) genotypes in type I and type II men, but not in women. A significant interaction for the DRD2 A1/A1 and 5-HTTLPR S+ polymorphisms was also found only in type I and type II men (Wang et al., 2014).

### **Pulmonary Hypertension (PPH)**

Smooth muscle cells (SMC) of pulmonary artery in PPH patients grow faster than controls when 5-HTT expression is stimulated by serotonin. As a result of these findings, 5-HTT has been shown to play a key role in the pathogenesis of SMC proliferation and 5-HTT polymorphisms are associated with PHH (Eddahibi et al., 2001). The higher 5-HTT expression was reported in pulmonary artery SMC in patients with L/L genotypes compared to L/S and S/S genotypes, resulting in more severe PPH. (Eddahibi et al., 2003). In a study on the association of idiopathic PPH with 5-HTT polymorphism in children, the L/L genotype has been reported to be associated with the aetiology of PPH (Vachharajani and Saunders, 2005).

## **References**

- Adamsen D, Meili D, Blau N, Thöny B, Ramaekers V. Autism associated with low 5-hydroxyindolacetic acid in CSF and the heterozygous SLC6A4 gene Gly56Ala plus 5-HTTLPR L/L promoter variants. *Mol Genet Metab.* 2011 Mar;102(3):368-73
- Ahmed BA, Bukhari IA, Jeffus BC, Harney JT, Thyparambil S, Ziu E, Fraer M, Rusch NJ, Zimniak P, Lupashin V, Tang D, Kilic F. The cellular distribution of serotonin transporter is impeded on serotonin-altered vimentin network. *PLoS One.* 2009;4(3):e4730
- Ahmed BA, Jeffus BC, Bukhari SI, Harney JT, Unal R, Lupashin VV, van der Sluijs P, Kilic F. Serotonin transamidates Rab4 and facilitates its binding to the C terminus of serotonin transporter. *J Biol Chem.* 2008 Apr 4;283(14):9388-98
- Albani D, Vittori A, Batelli S, Polito L, De Mauro S, Galimberti D, Scarpini E, Lovati C, Mariani C, Forloni G. Serotonin transporter gene polymorphic element 5-HTTLPR increases the risk of sporadic Parkinson's disease in Italy. *Eur Neurol.* 2009;62(2):120-3
- Allen NC, Bagade S, McQueen MB, Ioannidis JP, Kavvoura FK, Khoury MJ, Tanzi RE, Bertram L. Systematic meta-analyses and field synopsis of genetic association studies in schizophrenia: the SzGene database. *Nat Genet.* 2008 Jul;40(7):827-34



Ansorge MS, Zhou M, Lira A, Hen R, Gingrich JA. Early-life blockade of the 5-HT transporter alters emotional behavior in adult mice *Science* 2004 Oct

29;306(5697):879-81

Aoki J, Ikeda K, Murayama O, Yoshihara E, Ogai Y, Iwahashi K. The association between personality, pain threshold and a single nucleotide polymorphism (rs3813034) in the 3'-untranslated region of the serotonin transporter gene (SLC6A4) *J Clin Neurosci* 2010 May;17(5):574-8

Babineau V, Green CG, Jolicoeur-Martineau A, Bouvette-Turcot AA, Minde K, Sassi R, St-André M, Carrey N, Atkinson L, Kennedy JL, Lydon J, Steiner M, Gaudreau H, Levitan R, Meaney M, Wazana A; MAVAN project. Prenatal depression and 5-HTTLPR interact to predict dysregulation from 3 to 36 months--a differential susceptibility model *J Child Psychol Psychiatry* 2015 Jan;56(1):21-9

Baca-Garcia E, Vaquero-Lorenzo C, Diaz-Hernandez M, Rodriguez-Salgado B, Dolengevich-Segal H, Arrojo-Romero M, Botillo-Martin C, Ceverino A, Piqueras JF, Perez-Rodriguez MM, Saiz-Ruiz J. Association between obsessive-compulsive disorder and a variable number of tandem repeats polymorphism in intron 2 of the serotonin transporter gene *Prog Neuropsychopharmacol Biol Psychiatry* 2007 Mar 30;31(2):416-20

Borroni B, Grassi M, Agosti C, Archetti S, Costanzi C, Cornali C, Caltagirone C, Caimi L, Di Luca M, Padovani A. Cumulative effect of COMT and 5-HTTLPR polymorphisms and their interaction with disease severity and comorbidities on the risk of psychosis in Alzheimer disease *Am J Geriatr Psychiatry* 2006 Apr;14(4):343-51

Bradley SL, Dodelzon K, Sandhu HK, Philibert RA. Relationship of serotonin transporter gene polymorphisms and haplotypes to mRNA transcription *Am J Med Genet B Neuropsychiatr Genet* 2005 Jul 5;136B(1):58-61

Brenner B, Harney JT, Ahmed BA, Jeffus BC, Unal R, Mehta JL, Kilic F. Plasma serotonin levels and the platelet serotonin transporter *J Neurochem* 2007 Jul;102(1):206-15

Brown GL, Ebert MH, Goyer PF, Jimerson DC, Klein WJ, Bunney WE, Goodwin FK. Aggression, suicide, and serotonin: relationships to CSF amine metabolites *Am J Psychiatry* 1982 Jun;139(6):741-6

Camarena B, González L, Hernández S, Caballero A. SLC6A4 rare variant associated with eating disorders in Mexican patients *J Psychiatr Res* 2012 Aug;46(8):1106-7

Canli T, Lesch KP. Long story short: the serotonin transporter in emotion regulation and social cognition *Nat Neurosci* 2007 Sep;10(9):1103-9

Cao H, Gu H, Qiu W, Zuo W, Zheng L, Wang Z, Hu Z, Chen Y. Association study of serotonin transporter gene polymorphisms and ventricular septal defects related possible pulmonary arterial hypertension in Chinese population *Clin Exp Hypertens* 2009 Oct;31(7):605-14

Caspi A, Sugden K, Moffitt TE, Taylor A, Craig IW, Harrington H, McClay J, Mill J, Martin J, Braithwaite A, Poulton R. Influence of life stress on depression: moderation by a polymorphism in the 5-HTT gene *Science* 2003 Jul 18;301(5631):386-9

Catalano M. The challenges of psychopharmacogenetics *Am J Hum Genet* 1999 Sep;65(3):606-10

Chen NH, Reith ME, Quick MW. Synaptic uptake and beyond: the sodium- and chloride-dependent neurotransmitter transporter family SLC6 *Pflugers Arch* 2004 Feb;447(5):519-31

Cho HJ, Meira-Lima I, Cordeiro Q, Michelon L, Sham P, Vallada H, Collier DA. Population-based and family-based studies on the serotonin transporter gene polymorphisms and bipolar disorder: a systematic review and meta-analysis *Mol Psychiatry* 2005 Aug;10(8):771-81

Devlin AM, Brain U, Austin J, Oberlander TF. Prenatal exposure to maternal depressed mood and the MTHFR C677T variant affect SLC6A4 methylation in infants at birth *PLoS One* 2010 Aug 16;5(8):e12201

Eddahibi S, Humbert M, Fadel E, Raffestin B, Darmon M, Capron F, Simonneau G, Dartevelle P, Hamon M, Adnot S. Serotonin transporter overexpression is responsible for pulmonary artery smooth muscle hyperplasia in primary pulmonary hypertension *J Clin Invest* 2001 Oct;108(8):1141-50

Esterling LE, Yoshikawa T, Turner G, Badner JA, Bengel D, Gershon ES, Berrettini WH, Detera-Wadleigh SD. Serotonin transporter (5-HTT) gene and bipolar affective disorder *Am J Med Genet* 1998 Feb 7;81(1):37-40

Feinn R, Nellisery M, Kranzler HR. Meta-analysis of the association of a functional serotonin transporter promoter polymorphism with alcohol dependence *Am J Med Genet B Neuropsychiatr Genet* 2005 Feb 5;133B(1):79-84

Fuxman Bass JI, Sahni N, Shrestha S, Garcia-Gonzalez A, Mori A, Bhat N, Yi S, Hill DE, Vidal M, Walhout AJM. Human gene-centered transcription factor networks for enhancers and disease variants *Cell* 2015 Apr 23;161(3):661-673

Gaudet P, Livstone MS, Lewis SE, Thomas PD. Phylogenetic-based propagation of functional annotations within the Gene Ontology consortium *Brief Bioinform* 2011 Sep;12(5):449-62

Gelernter J, Kranzler H, Coccaro EF, Siever LJ, New AS. Serotonin transporter protein gene polymorphism and personality measures in African American and European American subjects *Am J Psychiatry* 1998 Oct;155(10):1332-8

Grabe HJ, Lange M, Wolff B, Völzke H, Lucht M, Freyberger HJ, John U, Cascorbi I. Mental and physical distress is modulated by a polymorphism in the 5-HT transporter gene interacting with social stressors and chronic disease burden *Mol Psychiatry* 2005 Feb;10(2):220-4

Green CG, Babineau V, Jolicoeur-Martineau A, Bouvette-Turcot AA, Minde K, Sassi R, St-André M, Carrey N, Atkinson L, Kennedy JL, Steiner M, Lydon J, Gaudreau H, Burack JA, Levitan R, Meaney MJ, Wazana A; Maternal Adversity, Vulnerability, and Neurodevelopment Research Team. Prenatal maternal depression and child serotonin transporter linked polymorphic region (5-HTTLPR) and dopamine receptor D4 (DRD4) genotype predict negative emotionality from 3 to 36 months *Dev Psychopathol* 2017 Aug;29(3):901-917

Gyawali S, Subaran R, Weissman MM, Hershkowitz D, McKenna MC, Talati A, Fyer AJ, Wickramaratne P, Adams PB, Hodge SE, Schmidt CJ, Bannon MJ, Glatt CE. Association of a polyadenylation polymorphism in the serotonin transporter and panic disorder *Biol Psychiatry* 2010 Feb 15;67(4):331-8

Hariri AR, Mattay VS, Tessitore A, Kolachana B, Fera F, Goldman D, Egan MF, Weinberger DR. Serotonin transporter genetic variation and the response of the human amygdala *Science* 2002 Jul 19;297(5580):400-3

Harrison SJ, Tyrer AE, Levitan RD, Xu X, Houle S, Wilson

AA, Nobrega JN, Rusjan PM, Meyer JH. Light therapy and serotonin transporter binding in the anterior cingulate and

- prefrontal cortex *Acta Psychiatr Scand* 2015 Nov;132(5):379-88
- Hartley CA, McKenna MC, Salman R, Holmes A, Casey BJ, Phelps EA, Glatt CE. Serotonin transporter polyadenylation polymorphism modulates the retention of fear extinction memory *Proc Natl Acad Sci U S A* 2012 Apr 3;109(14):5493-8
- Heils A, Teufel A, Petri S, Stöber G, Riederer P, Bengel D, Lesch KP. Allelic variation of human serotonin transporter gene expression *J Neurochem* 1996 Jun;66(6):2621-4
- Herman AI, Philbeck JW, Vasilopoulos NL, Depetrillo PB. Serotonin transporter promoter polymorphism and differences in alcohol consumption behaviour in a college student population *Alcohol Alcohol* 2003 Sep-Oct;38(5):446-9
- Hu XZ, Lipsky RH, Zhu G, Akhtar LA, Taubman J, Greenberg BD, Xu K, Arnold PD, Richter MA, Kennedy JL, Murphy DL, Goldman D. Serotonin transporter promoter gain-of-function genotypes are linked to obsessive-compulsive disorder *Am J Hum Genet* 2006 May;78(5):815-826
- Hung CF, Lung FW, Chen CH, O'Nions E, Hung TH, Chong MY, Wu CK, Wen JK, Lin PY. Association between suicide attempt and a tri-allelic functional polymorphism in serotonin transporter gene promoter in Chinese patients with schizophrenia *Neurosci Lett* 2011 Oct 31;504(3):242-6
- Kapur S, Remington G. Serotonin-dopamine interaction and its relevance to schizophrenia *Am J Psychiatry* 1996 Apr;153(4):466-76
- Kilic F, Murphy DL, Rudnick G. A human serotonin transporter mutation causes constitutive activation of transport activity *Mol Pharmacol* 2003 Aug;64(2):440-6
- Landaas ET, Johansson S, Jacobsen KK, Ribasés M, Bosch R, Sánchez-Mora C, Jacob CP, Boreatti-Hümmer A, Kreiker S, Lesch KP, Kiemenev LA, Kooij JJ, Kan C, Buitelaar JK, Faraone SV, Halmøy A, Ramos-Quiroga JA, Cormand B, Reif A, Franke B, Mick E, Knappskog PM, Haavik J. An international multicenter association study of the serotonin transporter gene in persistent ADHD *Genes Brain Behav* 2010 Jul;9(5):449-58
- Lasky-Su JA, Faraone SV, Glatt SJ, Tsuang MT. Meta-analysis of the association between two polymorphisms in the serotonin transporter gene and affective disorders *Am J Med Genet B Neuropsychiatr Genet* 2005 Feb 5;133B(1):110-5
- Lesch KP, Bengel D, Heils A, Sabol SZ, Greenberg BD, Petri S, Benjamin J, Müller CR, Hamer DH, Murphy DL. Association of anxiety-related traits with a polymorphism in the serotonin transporter gene regulatory region *Science* 1996 Nov 29;274(5292):1527-31
- Li J, Lin H, Zhu X, Li L, Wang X, Sun W, Wu X, Liu A, Niu F, Wang Y, Liu Y. Association study of functional polymorphisms in serotonin transporter gene with temporal lobe epilepsy in Han Chinese population *Eur J Neurol* 2012 Feb;19(2):351-3
- Li LB, Chen N, Ramamoorthy S, Chi L, Cui XN, Wang LC, Reith ME. The role of N-glycosylation in function and surface trafficking of the human dopamine transporter *J Biol Chem* 2004 May 14;279(20):21012-20
- Müller HK, Wiborg O, Haase J. Subcellular redistribution of the serotonin transporter by secretory carrier membrane protein 2 *J Biol Chem* 2006 Sep 29;281(39):28901-9
- MacKenzie A, Quinn J. A serotonin transporter gene intron 2 polymorphic region, correlated with affective disorders, has allele-dependent differential enhancer-like properties in the mouse embryo *Proc Natl Acad Sci U S A* 1999 Dec 21;96(26):15251-5
- MacLean MR, Deuchar GA, Hicks MN, Morecroft I, Shen S, Sheward J, Colston J, Loughlin L, Nilsen M, Dempsey Y, Harmar A. Overexpression of the 5-hydroxytryptamine transporter gene: effect on pulmonary hemodynamics and hypoxia-induced pulmonary hypertension *Circulation* 2004 May 4;109(17):2150-5
- Margolis KG, Li Z, Stevanovic K, Saurman V, Israelyan N, Anderson GM, Snyder I, Veenstra-VanderWeele J, Blakely RD, Gershon MD. Serotonin transporter variant drives preventable gastrointestinal abnormalities in development and function *J Clin Invest* 2016 Jun 1;126(6):2221-35
- Maron E, Shlik J. Serotonin function in panic disorder: important, but why? *Neuropsychopharmacology* 2006 Jan;31(1):1-11 Review
- Marziniak M, Mössner R, Schmitt A, Lesch KP, Sommer C. A functional serotonin transporter gene polymorphism is associated with migraine with aura *Neurology* 2005 Jan 11;64(1):157-9
- Melikian HE, Ramamoorthy S, Tate CG, Blakely RD. Inability to N-glycosylate the human norepinephrine transporter reduces protein stability, surface trafficking, and transport activity but not ligand recognition *Mol Pharmacol* 1996 Aug;50(2):266-76
- Mendlewicz J, Massat I, Souery D, Del-Favero J, Oruc L, Nöthen MM, Blackwood D, Muir W, Battersby S, Lerer B, Segman RH, Kaneva R, Serretti A, Lilli R, Lorenzi C, Jakovljevic M, Ivezic S, Rietschel M, Milanova V, Van Broeckhoven C. Serotonin transporter 5HTTLPR polymorphism and affective disorders: no evidence of association in a large European multicenter study *Eur J Hum Genet* 2004 May;12(5):377-82
- Mitra N, Sinha S, Ramya TN, Suroliya A. N-linked oligosaccharides as outfitters for glycoprotein folding, form and function *Trends Biochem Sci* 2006 Mar;31(3):156-63
- Moya PR, Wendland JR, Rubenstein LM, Timpano KR, Heiman GA, Tischfield JA, King RA, Andrews AM, Ramamoorthy S, McMahon FJ, Murphy DL. Common and rare alleles of the serotonin transporter gene, SLC6A4, associated with Tourette's disorder *Mov Disord* 2013 Aug;28(9):1263-70
- Munafò MR, Lingford-Hughes AR, Johnstone EC, Walton RT. Association between the serotonin transporter gene and alcohol consumption in social drinkers *Am J Med Genet B Neuropsychiatr Genet* 2005 May 5;135B(1):10-4
- Murphy DL, Fox MA, Timpano KR, Moya PR, Ren-Patterson R, Andrews AM, Holmes A, Lesch KP, Wendland JR. How the serotonin story is being rewritten by new gene-based discoveries principally related to SLC6A4, the serotonin transporter gene, which functions to influence all cellular serotonin systems *Neuropharmacology* 2008 Nov;55(6):932-60
- Narita N, Narita M, Takashima S, Nakayama M, Nagai T, Okado N. Serotonin transporter gene variation is a risk factor for sudden infant death syndrome in the Japanese population *Pediatrics* 2001 Apr;107(4):690-2
- Oberlander TF, Papsdorf M, Brain UM, Misri S, Ross C, Grunau RE. Prenatal effects of selective serotonin reuptake inhibitor antidepressants, serotonin transporter promoter genotype (SLC6A4), and maternal mood on child behavior at 3 years of age *Arch Pediatr Adolesc Med* 2010 May;164(5):444-51

- Ogilvie AD, Battersby S, Bubbs VJ, Fink G, Harmar AJ, Goodwin GM, Smith CA. Polymorphism in serotonin transporter gene associated with susceptibility to major depression *Lancet* 1996 Mar 16;347(9003):731-3
- Ohara K, Nagai M, Tsukamoto T, Tani K, Suzuki Y, Ohara K. Functional polymorphism in the serotonin transporter promoter at the SLC6A4 locus and mood disorders *Biol Psychiatry* 1998 Oct 1;44(7):550-4
- Olivares L, Aragón C, Giménez C, Zafra F. The role of N-glycosylation in the targeting and activity of the GLYT1 glycine transporter *J Biol Chem* 1995 Apr 21;270(16):9437-42
- Oppenheimer CW, Hankin BL, Young JF, Smolen A. Youth genetic vulnerability to maternal depressive symptoms: 5-HTTLPR as moderator of intergenerational transmission effects in a multiwave prospective study *Depress Anxiety* 2013 Mar;30(3):190-6
- Ozaki N, Goldman D, Kaye WH, Plotnicov K, Greenberg BD, Lappalainen J, Rudnick G, Murphy DL. Serotonin transporter missense mutation associated with a complex neuropsychiatric phenotype *Mol Psychiatry* 2003 Nov;8(11):933-6
- Page DT, Kuti OJ, Prestia C, Sur M. Haploinsufficiency for Pten and Serotonin transporter cooperatively influences brain size and social behavior *Proc Natl Acad Sci U S A* 2009 Feb 10;106(6):1989-94
- Palmer AM, Stratmann GC, Procter AW, Bowen DM. Possible neurotransmitter basis of behavioral changes in Alzheimer's disease *Ann Neurol* 1988 Jun;23(6):616-20
- Perroud N, Salzmann A, Saiz PA, Baca-Garcia E, Sarchiapone M, Garcia-Portilla MP, Carli V, Vaquero-Lorenzo C, Jaussent I, Mouton D, Vessaz M, Huguelet P, Courtet P, Malafosse A; European Research Consortium for Suicide (EURECA). Rare genotype combination of the serotonin transporter gene associated with treatment response in severe personality disorder *Am J Med Genet B Neuropsychiatr Genet* 2010 Dec 5;153B(8):1494-7
- Petito A, Altamura M, Iuso S, Padalino FA, Sessa F, D'Andrea G, Margaglione M, Bellomo A. The Relationship between Personality Traits, the 5HTT Polymorphisms, and the Occurrence of Anxiety and Depressive Symptoms in Elite Athletes *PLoS One* 2016 Jun 3;11(6):e0156601
- Prasad HC, Steiner JA, Sutcliffe JS, Blakely RD. Enhanced activity of human serotonin transporter variants associated with autism *Philos Trans R Soc Lond B Biol Sci* 2009 Jan 27;364(1514):163-73
- Prasad HC, Zhu CB, McCauley JL, Samuvel DJ, Ramamoorthy S, Shelton RC, Hewlett WA, Sutcliffe JS, Blakely RD. Human serotonin transporter variants display altered sensitivity to protein kinase G and p38 mitogen-activated protein kinase *Proc Natl Acad Sci U S A* 2005 Aug 9;102(32):11545-50
- Ramamoorthy S, Blakely RD. Phosphorylation and sequestration of serotonin transporters differentially modulated by psychostimulants *Science* 1999 Jul 30;285(5428):763-6
- Ramos N, Reichert JG, Corwin TE, Smith CJ, Silverman JM, Hollander E, Buxbaum JD. Lack of evidence for association of the serotonin transporter gene SLC6A4 with autism *Biol Psychiatry* 2006 Jul 15;60(2):186-91
- Rao S, Leung CS, Lam MH, Wing YK, Waye MM, Tsui SK. Resequencing three candidate genes discovers seven potentially deleterious variants susceptibility to major depressive disorder and suicide attempts in Chinese *Gene* 2017 Mar 1;603:34-41
- Rasmussen TN, Plenge P, Bay T, Egebjerg J, Gether U. A single nucleotide polymorphism in the human serotonin transporter introduces a new site for N-linked glycosylation *Neuropharmacology* 2009 Sep;57(3):287-94
- Rotondo A, Mazzanti C, Dell'Osso L, Rucci P, Sullivan P, Bouanani S, Gonnelli C, Goldman D, Cassano GB. Catechol o-methyltransferase, serotonin transporter, and tryptophan hydroxylase gene polymorphisms in bipolar disorder patients with and without comorbid panic disorder *Am J Psychiatry* 2002 Jan;159(1):23-9
- Saiz PA, Garcia-Portilla MP, Arango C, Morales B, Bascaran MT, Martinez-Barrondo S, Florez G, Sotomayor E, Paredes B, Alvarez C, San Narciso G, Carreño E, Bombin I, Alvarez V, Coto E, Fernandez JM, Boussoño M, Bobes J. Association study between obsessive-compulsive disorder and serotonergic candidate genes *Prog Neuropsychopharmacol Biol Psychiatry* 2008 Apr 1;32(3):765-70
- Samuvel DJ, Jayanthi LD, Bhat NR, Ramamoorthy S. A role for p38 mitogen-activated protein kinase in the regulation of the serotonin transporter: evidence for distinct cellular mechanisms involved in transporter surface expression *J Neurosci* 2005 Jan 5;25(1):29-41
- Sanjuan J, Martin-Santos R, Garcia-Estevé L, Carot JM, Guíllamat R, Gutierrez-Zotes A, Gornemann I, Canellas F, Baca-Garcia E, Jover M, Navines R, Valles V, Vilella E, de Diego Y, Castro JA, Ivorra JL, Gelabert E, Guitart M, Labad A, Mayoral F, Roca M, Gratacos M, Costas J, van Os J, de Frutos R. Mood changes after delivery: role of the serotonin transporter gene *Br J Psychiatry* 2008 Nov;193(5):383-8
- Scott M, Reading HW, Loudon JB. Studies on human blood platelets in affective disorder *Psychopharmacology (Berl)* 1979 Jan 31;60(2):131-5
- Sen S, Burmeister M, Ghosh D. Meta-analysis of the association between a serotonin transporter promoter polymorphism (5-HTTLPR) and anxiety-related personality traits *Am J Med Genet B Neuropsychiatr Genet* 2004 May 15;127B(1):85-9
- Seneviratne C, Huang W, Ait-Daoud N, Li MD, Johnson BA. Characterization of a functional polymorphism in the 3' UTR of SLC6A4 and its association with drinking intensity *Alcohol Clin Exp Res* 2009 Feb;33(2):332-9
- Strug LJ, Suresh R, Fyer AJ, Talati A, Adams PB, Li W, Hodge SE, Gilliam TC, Weissman MM. Panic disorder is associated with the serotonin transporter gene (SLC6A4) but not the promoter region (5-HTTLPR) *Mol Psychiatry* 2010 Feb;15(2):166-76
- Sugawara H, Iwamoto K, Bundo M, Ueda J, Miyauchi T, Komori A, Kazuno A, Adati N, Kusumi I, Okazaki Y, Ishigooka J, Kojima T, Kato T. Hypermethylation of serotonin transporter gene in bipolar disorder detected by epigenome analysis of discordant monozygotic twins *Transl Psychiatry* 2011 Jul 26;1:e24
- Sukonick DL, Pollock BG, Sweet RA, Mulsant BH, Rosen J, Klunk WE, Kastango KB, DeKosky ST, Ferrell RE. The 5-HTTLPR\*<sup>S</sup>/L polymorphism and aggressive behavior in Alzheimer disease *Arch Neurol* 2001 Sep;58(9):1425-8
- Sutcliffe JS, Delahanty RJ, Prasad HC, McCauley JL, Han Q, Jiang L, Li C, Folstein SE, Blakely RD. Allelic heterogeneity at the serotonin transporter locus (SLC6A4) confers susceptibility to autism and rigid-compulsive behaviors *Am J Hum Genet* 2005 Aug;77(2):265-79
- Tate CG, Blakely RD. The effect of N-linked glycosylation on activity of the Na(+)- and Cl(-)-dependent serotonin transporter expressed using recombinant baculovirus in insect cells *J Biol Chem* 1994 Oct 21;269(42):26303-10

- Taylor S. Molecular genetics of obsessive-compulsive disorder: a comprehensive meta-analysis of genetic association studies *Mol Psychiatry* 2013 Jul;18(7):799-805
- Tyrer AE, Levitan RD, Houle S, Wilson AA, Nobrega JN, Meyer JH. Increased Seasonal Variation in Serotonin Transporter Binding in Seasonal Affective Disorder *Neuropsychopharmacology* 2016 Sep;41(10):2447-54
- Vachharajani A, Saunders S. Allelic variation in the serotonin transporter (5HTT) gene contributes to idiopathic pulmonary hypertension in children *Biochem Biophys Res Commun* 2005 Aug 26;334(2):376-9
- Vallender EJ, Priddy CM, Hakim S, Yang H, Chen GL, Miller GM. Functional variation in the 3' untranslated region of the serotonin transporter in human and rhesus macaque *Genes Brain Behav* 2008 Aug;7(6):690-7
- Wang TY, Lee SY, Chen SL, Chang YH, Chen SH, Huang SY, Tzeng NS, Wang CL, Yeh PH, Chen KC, Lee IH, Yeh TL, Yang YK, Lu RB. Gender-specific association of the SLC6A4 and DRD2 gene variants in bipolar disorder *Int J Neuropsychopharmacol* 2014 Feb;17(2):211-22
- Wankerl M, Miller R, Kirschbaum C, Hennig J, Stalder T, Alexander N. Effects of genetic and early environmental risk factors for depression on serotonin transporter expression and methylation profiles *Transl Psychiatry* 2014 Jun 17;4:e402
- Weese-Mayer DE, Berry-Kravis EM, Maher BS, Silvestri JM, Curran ME, Marazita ML. Sudden infant death syndrome: association with a promoter polymorphism of the serotonin transporter gene *Am J Med Genet A* 2003 Mar 15;117A(3):268-74
- Wendland JR, Moya PR, Kruse MR, Ren-Patterson RF, Jensen CL, Timpano KR, Murphy DL. A novel, putative gain-of-function haplotype at SLC6A4 associates with obsessive-compulsive disorder *Hum Mol Genet* 2008 Mar 1;17(5):717-23
- Willeit M, Praschak-Rieder N, Neumeister A, Zill P, Leisch F, Stastny J, Hilger E, Thierry N, Konstantinidis A, Winkler D, Fuchs K, Sieghart W, Aschauer H, Ackenheil M, Bondy B, Kasper S. A polymorphism (5-HTTLPR) in the serotonin transporter promoter gene is associated with DSM-IV depression subtypes in seasonal affective disorder *Mol Psychiatry* 2003 Nov;8(11):942-6
- Yamashita A, Singh SK, Kawate T, Jin Y, Gouaux E. Crystal structure of a bacterial homologue of Na<sup>+</sup>/Cl<sup>-</sup>-dependent neurotransmitter transporters *Nature* 2005 Sep 8;437(7056):215-23
- Yoshida T, Hojo S, Sekine S, Sawada S, Okumura T, Nagata T, Shimada Y, Tsukada K. Expression of aquaporin-1 is a poor prognostic factor for stage II and III colon cancer *Mol Clin Oncol* 2013 Nov;1(6):953-958
- Zhang Y, Bertolino A, Fazio L, Blasi G, Rampino A, Romano R, Lee ML, Xiao T, Papp A, Wang D, Sadée W. Polymorphisms in human dopamine D2 receptor gene affect gene expression, splicing, and neuronal activity during working memory *Proc Natl Acad Sci U S A* 2007 Dec 18;104(51):20552-7
- Zhu CB, Hewlett WA, Feoktistov I, Biaggioni I, Blakely RD. Adenosine receptor, protein kinase G, and p38 mitogen-activated protein kinase-dependent up-regulation of serotonin transporters involves both transporter trafficking and activation *Mol Pharmacol* 2004 Jun;65(6):1462-74

---

*This article should be referenced as such:*

Gurbanov R, Kalkanci B. SLC6A4 (solute carrier family 6 member 4). *Atlas Genet Cytogenet Oncol Haematol*. 2020; 24(1):39-50.

---